

Appendix B

CHART Assessment for the Lower Columbia River Chinook Salmon ESU

CHART Participants

The CHART for this ESU consisted of the following NOAA Fisheries biologists: Ben Meyer (CHART Leader), Michelle Day, Patty Dornbusch, Dan Guy, Lynne Krasnow, Lance Kruzic, Nancy Munn, Mindy Simmons, Cathy Tortorici, and Rich Turner. This CHART assessment also benefitted from review and comments from the Oregon and Washington Departments of Fish and Wildlife (ODFW).

ESU Description

The Lower Columbia River Chinook ESU was listed as a threatened species in 1999. The ESU includes all naturally spawned populations of Chinook salmon from the Columbia River and its tributaries from its mouth at the Pacific Ocean upstream to a transitional point between Washington and Oregon east of the Hood River and the White Salmon River, and includes the Willamette River to Willamette Falls, Oregon, exclusive of spring-run Chinook salmon in the Clackamas River (64 FR 14308; March 24, 1999). The agency recently conducted a review to update the ESU's status, taking into account new information and considering the net contribution of artificial propagation efforts in the ESU. We recently published the results of this review and concluded that Lower Columbia River Chinook salmon (including 17 hatchery programs) should remain listed as threatened (70 FR 37160; June 28, 2005).

The following brief description is based largely on life history information and excerpts from the report of the Lower Columbia Fish Recovery Board (LCFRB 2003) and the Willamette/Lower Columbia River Technical Recovery Team's (TRT) recent review of historical population structure for this ESU (Myers et al. 2003).

Of the Pacific salmon, Chinook salmon exhibit the most diverse and complex life history strategies. Chinook salmon follow one of two general freshwater cycles: stream or ocean type. After emerging from the gravel, stream-type Chinook salmon reside in fresh water for a year or more before migrating to the ocean. Ocean-type Chinook salmon migrate to the ocean within their first year. These two types of Chinook salmon have different life history traits, geographic distribution, and genetic characteristics. Chinook in the lower Columbia River generally follow an ocean-type life history cycle.

Runs are designated on the basis of when adults enter freshwater; however, distinct runs may also differ in the degree of maturation at river entry and time of spawning. Early,

spring-run (stream-maturing) Chinook salmon tend to enter freshwater as immature or bright fish, migrate upriver (holding in suitable thermal refuges for several months), and finally spawn in late summer and early autumn. Late, fall-run (ocean maturing) Chinook salmon enter freshwater at an advanced stage of maturity, move rapidly to their spawning areas on the main stem or lower tributaries of the rivers, and spawn within a few days or weeks of freshwater entry. Fall Chinook dominate Chinook salmon runs in this ESU. Today, the once abundant natural runs of fall and spring Chinook have been largely replaced by hatchery production. Large Chinook runs continue to return to many of their natal streams, but there are few sustained native, naturally reproducing populations.

Adult spring Chinook return to the Columbia River at 4 to 5 years of age. They enter the Columbia River in March and April and generally enter natal basins from March through June, well in advance of spawning in August and September. Spring Chinook typically spawn in headwater areas where higher gradient habitat exists. Successful spawning depends on sufficient clean gravel of the right size, in addition to the constant need of adequate flows and water quality. Fall Chinook return to the Columbia River at 3 to 4 years of age, although 5-year olds are common in some populations. They enter fresh water from August to September and spawning generally occurs from late September to November, with peak spawning activity in mid-October. Bright fall Chinook adults enter the Columbia River August to October; dominant age class varies by population and brood year, but is typically age 4. Spawning occurs in November to January, with peak spawning in mid November.

Chinook salmon eggs incubate throughout the autumn and winter months. As with other salmonids, water temperature controls incubation time and affects survival. During incubation, clean, well-oxygenated water flow is critical. Floods/scouring, dewatering, and sedimentation can result in high egg mortality. In the lower Columbia River, spring Chinook fry emerge from the gravel from November through March; peak emergence time is likely December and January. Fall Chinook fry generally emerge from the gravel in April, depending on the time of egg deposition and incubation water temperature. The emerging fry quickly migrate to quiet waters and off-stream areas where they can find food and protection from predators.

After emerging from the gravel in the spring, most fall Chinook fry rear in the freshwater habitat for 1 to 4 months before emigrating to the ocean as subyearlings. A few fall Chinook remain in fresh water until their second spring and emigrate as yearlings. Conversely, spring Chinook emerge from the gravel earlier than fall Chinook, generally in the late winter/early spring. Normally, spring Chinook spend one full year in fresh water and emigrate to sea in their second spring. After emergence fry generally search

for suitable rearing habitat within side sloughs, side channels, spring-fed seep areas and along the outer edges of the stream. These quiet-water side margin and off-channel slough areas are vital for early juvenile habitat. The presence of woody debris and overhead cover aid in food and nutrient inputs, and provide protection from predators during early freshwater residence.

Juvenile Chinook salmon in freshwater feed on a variety of terrestrial and aquatic insects and crustaceans, while subadults feed on similar items as well as larger prey including fishes, shrimp, and squid (Scott and Crossman, 1973). One study noted that adults in marine waters forage on a large array of fish species, especially herring and sand lance (Pritchard and Tester 1944 as cited in Scott and Crossman 1973).

Recovery Planning Status

The Willamette/Lower Columbia TRT identified 31 historical demographically independent Chinook salmon populations in this ESU (Myers et al. 2003). It is estimated that eight to ten historical populations in the ESU have been extirpated or nearly so. The TRT has grouped populations within the ESU into three life-history types (spring-, fall-, and late fall-run) and three ecological spawning zones (Coast Range, Cascade, and Columbia Gorge) (McElhany et al. 2002). Recovery planning will likely emphasize the need for a geographical distribution of viable populations across the range of life-history types and ecological zones (Ruckelshaus et al. 2002, McElhany et al. 2003). A draft recovery plan for the Washington management unit of this ESU was completed by the Lower Columbia Fish Recovery Board (LCFRB 2004) and released by NMFS for public comment in April 2005. NMFS expects to use this plan as an interim regional recovery plan until a plan for the whole ESU is completed. A preliminary draft plan for Oregon areas of the ESU is expected by the end of 2005. The CHART considered the LCFRB plan and the TRT products in rating each habitat area, but did not have the benefit of regional recovery plans throughout the range of this ESU. We anticipate that, as recovery planning proceeds, we will have better information and may revise our recommendations regarding critical habitat designation.

CHART Area Assessments

The CHART assessment for this ESU addressed 10 subbasins containing 47 occupied watersheds, as well as the lower Columbia River rearing/migration corridor. As part of its assessment the CHART considered the conservation value of each watershed in the context of the populations within the strata identified by the TRT (McElhany et al. 2002). Information is presented below by USGS subbasin because they present a convenient and

systematic way to organize the CHART's watershed assessments for this ESU and their names are generally more recognizable because they typically identify major river systems.

Middle Columbia/Hood Subbasin (HUC4# 17070105)

The Middle Columbia/Hood subbasin is located in the eastern portion of the Columbia River gorge of Oregon and Washington. Occupied watersheds in this subbasin are contained in Hood River, Multnomah, and Wasco counties in Oregon, and Klickitat and Skamania counties in Washington. The subbasin contains 13 watersheds, 8 of which are occupied by this ESU. Occupied watersheds encompass approximately 1,370 mi² and 1,494 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) and Washington Department of Fish and Wildlife (WDFW) identify approximately 145 miles of occupied riverine habitat in the watersheds, including a 23-mile segment of the Columbia River (ODFW 2003a,b; WDFW 2003). Myers et al. (2003) identified a single ecological zone (Columbia Gorge) containing four fall-run (Lower Gorge tributaries, Upper Gorge tributaries, Big White Salmon River, and Hood River) and two spring-run (Big White Salmon River and Hood River) historical demographically independent populations in this subbasin. The Upper Gorge tributaries fall-run and Big White Salmon fall- and spring-run populations have been classified by the TRT as "core" populations, i.e., historically abundant and "may offer the most likely path to recovery" (McElhany et al. 2003). Native spring-run Chinook salmon are believed to be extirpated in this subbasin, although efforts are underway to reestablish these fish.

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B1 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin were of either high or medium conservation value to the ESU. Of the eight HUC5s reviewed, six were rated as having high and two were rated as having medium conservation value. The CHART noted that two HUC5s (Middle Columbia/Eagle Creek and Middle Columbia/Grays Creek) contain a high value rearing and migration corridor in the Columbia River connecting high value upstream watersheds with downstream reaches and the ocean. Table B2 summarizes the

CHART's PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

The CHART also considered whether blocked historical habitats above Condit Dam (on the White Salmon River) may be essential for conservation of the ESU. The Team determined that accessing this habitat would likely provide a benefit to the ESU, especially for spring-run Chinook salmon of which there are only two historical populations in the Gorge region. However, the CHART concluded that it was unclear whether the areas above Condit Dam are essential for conservation of the entire ESU, especially in comparison to other, more extensive, historical habitats that may be of greater potential benefit to the ESU (e.g., areas in the Upper Lewis River).

Lower Columbia/Sandy Subbasin (HUC4# 17080001)

The Lower Columbia/Sandy subbasin is located in the western portion of the Columbia River gorge of Oregon and Washington. Occupied watersheds in this subbasin are contained in Clackamas, Columbia, and Multnomah counties in Oregon, and Clark and Skamania counties in Washington. The subbasin contains nine watersheds, all of which are occupied by this ESU. Occupied watersheds encompass approximately 1,076 mi² and 1,316 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) and Washington Department of Fish and Wildlife (WDFW) identify approximately 217 miles of occupied riverine habitat in the watersheds, including a 26-mile segment of the Columbia River (ODFW 2003a,b; WDFW 2003). Myers et al. (2003) identified two ecological zones (Cascade and Columbia Gorge) containing five fall-run (Lower Gorge tributaries, Sandy River early fall, Sandy River late fall, Washougal River, and Salmon Creek/Lewis River) and one spring-run (Sandy River) historical demographically independent populations in this subbasin. The Sandy River late fall- and spring-run Chinook salmon have been classified by the TRT as “core” populations, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003). Also, the TRT classified the Sandy River spring- and late fall-runs and the Salmon Creek/Lewis River fall-run as genetic legacy populations, i.e., some of “the most intact representatives of the genetic character of the ESU” (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in

the watersheds. Map B2 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin ranged from high to low conservation value to the ESU. Of the nine HUC5s reviewed, seven were rated as having high, one was rated as having medium, and one was rated as having low conservation value. The CHART also noted that one HUC5 (Columbia Gorge Tributaries) contains a high value rearing and migration corridor in the Columbia River connecting high value upstream watersheds with downstream reaches and the ocean. Table B2 summarizes the CHART's PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

The CHART also concluded that inaccessible reaches above the Bull Run Dam complex in the Bull Run River HUC5 may be essential to the conservation of the ESU. The CHART concluded that these unoccupied areas may be essential because (1) they once supported TRT core and genetic legacy populations (Sandy River spring- and late fall-runs) and (2) they contain non-inundated habitats that are likely in good to excellent condition (i.e., the watershed provides domestic drinking water for the City of Portland and may have been some of the better spawning areas) (Sieglitz 2002, McElhany et al. 2003). The CHART noted that NOAA Fisheries' status review of this ESU stated that habitat loss due to "extensive hydropower development projects" posed a serious threat to this ESU (NOAA Fisheries 2003). This report also expressed serious concerns associated with dramatic declines in the spring-run life history type (which inhabits this watershed). Therefore, the CHART concluded that the ESU would likely benefit if the extant population of spring-run fish had access to spawning/rearing habitat upstream and that these areas may warrant consideration as critical habitat. However, the Team also considered historical areas in the upper Lewis River basin (see below) to have greater conservation potential than areas above the Bull Run Dam complex.

Lewis Subbasin (HUC4# 17080002)

The Lewis subbasin is located in southwest Washington and contained in Clark, Cowlitz, and Skamania counties (a very small and unoccupied portion in the uppermost watershed is contained in Yakima County). The subbasin contains six watersheds, two of which are currently occupied by this ESU and the remaining four are now blocked by Merwin Dam and others upstream. Occupied watersheds encompass approximately 456 mi² and 561 miles of streams. Fish distribution and habitat use data from the Washington Department of Fish and Wildlife (WDFW) identify approximately 68 miles of occupied riverine habitat in the watersheds (WDFW 2003). Myers et al. (2003) identified a single ecological zone (Cascade) containing one spring-run (Lewis River), one fall-run (Salmon

Creek/Lewis River) and one late fall-run (Lewis River) historical demographically independent populations in this subbasin. The TRT has classified the Lewis River spring- and late fall-run populations as “core” populations (historically abundant and “may offer the most likely path to recovery”) and the Lewis River late fall-run and Salmon Creek/Lewis River fall-run populations as genetic legacy populations (some of “the most intact representatives of the genetic character of the ESU”) (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B3 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that both of the occupied HUC5 watersheds in this subbasin were of high conservation value to the ESU. Table B2 summarizes the CHART’s PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

The CHART also concluded that inaccessible reaches above Merwin, Yale and Swift dams may be essential to the conservation of the ESU. The CHART believed that these unoccupied areas may be essential because (1) they once supported TRT core and genetic legacy populations and (2) they contain non-inundated habitats that are likely in good condition relative to other more urbanized watersheds in the Cascade region (Lower Columbia River Fish Recovery Board 2003, McElhany et al. 2003). The CHART noted that NOAA Fisheries’ status review of this ESU stated that habitat loss due to “extensive hydropower development projects” posed a serious threat to this ESU (NOAA Fisheries 2003). This report also expressed serious concerns associated with dramatic declines in the spring-run life history type (which inhabits this watershed). Therefore, the CHART concluded that the ESU would likely benefit if the extant population of spring-run fish had access to spawning/rearing habitat upstream and that these areas may warrant consideration as critical habitat.

Lower Columbia/Clatskanie Subbasin (HUC4# 17080003)

The Lower Columbia/Clatskanie subbasin is located in southwest Washington and northwest Oregon. Occupied watersheds in this subbasin are contained in Clatsop and Columbia counties in Oregon, and Cowlitz, Lewis, Skamania, and Wahkiakum counties in Washington. The subbasin contains six watersheds, all of which are occupied by this

ESU. Occupied watersheds encompass approximately 841 mi² and 977 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) and Washington Department of Fish and Wildlife (WDFW) identify approximately 168 miles of occupied riverine habitat in the watersheds (ODFW 2003a,b; WDFW 2003). Myers et al. (2003) identified two ecological zones (Coast Range and Cascade) containing five fall-run (Elochoman River, Mill Creek, Kalama River, Clatskanie River, and Scappoose River) and one spring-run (Kalama River) historical demographically independent populations in this subbasin. The Elochoman River fall-run population has been classified by the TRT as a “core” population, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B4 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin ranged from high to low conservation value to the ESU. Of the six HUC5s reviewed, two were rated as having high, three were rated as having medium conservation value, and one was rated as having low conservation value to the ESU. Table B2 summarizes the CHART’s PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Upper Cowlitz Subbasin (HUC4# 17080004)

The Upper Cowlitz subbasin is located in southwest Washington and contained in Lewis, Pierce, Skamania, and Yakima counties. The subbasin contains five watersheds, all of which are occupied by this ESU. Occupied watersheds encompass approximately 1,030 mi² and 1,282 miles of streams. Fish distribution and habitat use data from the Washington Department of Fish and Wildlife (WDFW) identify approximately 104 miles of occupied riverine habitat in the watersheds (WDFW 2003). All of this habitat is located upstream of impassable dams (Mayfield and Mossyrock) and only accessible to anadromous fish via trap and haul operations. Myers et al. (2003) identified one ecological zone (Cascade) containing one fall-run (Upper Cowlitz River) and two spring-run (Upper Cowlitz River and Cispus River) historical demographically independent populations in this subbasin. Both spring-run populations have been classified by the TRT as “core” populations, i.e., historically abundant and “may offer the most likely path

to recovery” (McElhany et al. 2003). In addition, the TRT classified the Upper Cowlitz River spring-run population as a genetic legacy population, i.e., one of “the most intact representatives of the genetic character of the ESU.” However, there are significant uncertainties about the remaining stock structure in this subbasin (Myers et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B5 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin were all of high conservation value to the ESU. Table B2 summarizes the CHART’s PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Lower Cowlitz Subbasin (HUC4# 17080005)

The Lower Cowlitz subbasin is located in southwest Washington and contained in Cowlitz, Lewis, and Skamania counties. The subbasin contains eight watersheds, all of which are occupied by this ESU. Occupied watersheds encompass approximately 1,460 mi² and 1,510 miles of streams. Fish distribution and habitat use data from the Washington Department of Fish and Wildlife (WDFW) identify approximately 350 miles of occupied riverine habitat in the (WDFW 2003). Habitat in two HUC5 watersheds – Tilton River and Riffe Reservoir – is located upstream of impassable dams (Mayfield and Mossyrock) and only accessible to anadromous fish via trap and haul operations. Data from WDFW identified very little Chinook salmon distribution in the Riffe Reservoir HUC5 watershed (and did not identify the Riffe and Mayfield lakes as occupied habitat). However, the CHART determined that these lakes are occupied and contain PCEs for rearing/migrating juveniles based on information regarding migrants described in Wade (2000) as well as their own knowledge of trap and haul operations in this subbasin. Myers et al. (2003) identified one ecological zone (Cascade) containing four fall-run (Coweeman River, Toutle River, Lower Cowlitz River, and Upper Cowlitz River) and four spring-run (Toutle River, Tilton River, Upper Cowlitz River, and Cispus River) historical demographically independent populations in this subbasin. The latter two spring-run populations as well as the Toutle River and Lower Cowlitz River fall-run populations have been classified by the TRT as “core” populations, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003). In

addition, the TRT classified the Upper Cowlitz River spring-run and Coweeman River fall-run as genetic legacy populations, i.e., some of “the most intact representatives of the genetic character of the ESU.”

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B6 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART determined that the occupied HUC5 watersheds in this subbasin were of high or medium conservation value to the ESU. Of the eight HUC5s reviewed, four were rated as having high and four were rated as having medium conservation value to the ESU. The CHART also noted that four HUC5s (Riffe Reservoir, Jackson Prairie, East Willapa, and Coweeman River) contained high value rearing and migration corridors connecting high value upstream watersheds with downstream reaches and the ocean. Table B2 summarizes the CHART’s PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Lower Columbia Subbasin (HUC4# 17080006)

The Lower Columbia subbasin is located at the mouth of the Columbia River in southwest Washington and Northwest Oregon. Occupied watersheds in this subbasin are contained in Clatsop County, Oregon, and Lewis, Pacific, and Wahkiakum counties in Washington. The subbasin contains three watersheds, all of which are occupied by this ESU. Occupied watersheds encompass approximately 515 mi² and 638 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) and Washington Department of Fish and Wildlife (WDFW) identify approximately 122 miles of occupied riverine habitat in the watersheds (ODFW 2003a,b; WDFW 2003). Myers et al. (2003) identified a single ecological zone (Coast Range) containing three fall-run historical demographically independent populations in this subbasin (Grays River, Youngs Bay, and Big Creek). The Big Creek fall-run population has been classified by the TRT as a “core” population, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning,

rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B7 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin were of either high (Big Creek and Grays Bay) or medium (Youngs River) conservation value to the ESU. Table B2 summarizes the CHART's PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Middle Willamette Subbasin (HUC4# 17090007)

The portion of the Middle Willamette River subbasin occupied by this ESU is downstream of Willamette Falls and includes a single HUC5 watershed (Abernethy Creek) as well as a short segment (approximately 1 mile) of the Willamette River downstream of Willamette Falls. Occupied portions of this subbasin within the ESU's range are contained in Clackamas County, Oregon. The Abernethy Creek watershed encompasses approximately 134 mi² and 171 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) identify approximately 3 miles of occupied riverine habitat in the subbasin (ODFW 2003a,b). The occupied portions of the subbasin are in the Cascade ecological zone identified by Myers et al. (2003), but the TRT did not associate fish in this area with a historical demographically independent population (McElhany et al. 2003). However, the mouth of Abernethy Creek enters the Willamette upstream and in close proximity (less than 0.6 miles) to the mouth of the Clackamas River which does contain a fall-run population identified by the TRT.

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in the Abernethy Creek watershed contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B8 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the Abernethy Creek HUC5 watershed was of low conservation value to the ESU. Table B2 summarizes the CHART's PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Clackamas Subbasin (HUC4# 17090011)

The Clackamas subbasin is a Cascade Range drainage of the lower Willamette River and is contained in Clackamas and Marion counties, Oregon. The subbasin contains six watersheds, two of which are occupied by this ESU (Lower Clackamas and Eagle Creek). Occupied watersheds encompass approximately 270 mi² and 339 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) identify approximately 55 miles of occupied riverine habitat in the watersheds (ODFW 2003a,b). Myers et al. (2003) identified a single ecological zone (Cascade) containing a single historical demographically independent population in this subbasin (Clackamas River fall-run). This fall-run population has been classified by the TRT as a “core” population, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B9 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin were of high (Lower Clackamas River) and low (Eagle Creek) conservation value to the ESU. Table B2 summarizes the CHART’s PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Lower Willamette Subbasin (HUC4# 17090012)

The Lower Willamette subbasin is located at the confluence of the Willamette and Columbia rivers in Northwest Oregon. Occupied watersheds in this subbasin are contained in Clackamas, Multnomah, and Washington counties, Oregon. The subbasin contains three watersheds, all of which are occupied by this ESU. Occupied watersheds encompass approximately 407 mi² and 448 miles of streams. Fish distribution and habitat use data from the Oregon Department of Fish and Wildlife (ODFW) identify approximately 88 miles of occupied riverine habitat in the watersheds (ODFW 2003a,b). Myers et al. (2003) identified a single ecological zone (Cascade) containing two fall-run historical demographically independent populations in this subbasin (Clackamas River and Scappoose River). The Clackamas River fall-run population has been classified by the TRT as a “core” population, i.e., historically abundant and “may offer the most likely path to recovery” (McElhany et al. 2003).

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that all of the occupied areas in this subbasin contain one or more PCEs for this ESU. Table B1 summarizes the total number of occupied reaches identified for each HUC5 watershed as containing spawning, rearing, or migration PCEs, as well as management activities that may affect the PCEs in the watersheds. Map B10 depicts the specific areas in this subbasin occupied by the ESU and under consideration for critical habitat designation. The CHART also determined that the occupied HUC5 watersheds in this subbasin were of either high (Columbia Slough/Willamette) or medium (Johnson Creek and Scappoose Creek) conservation value to the ESU. The CHART also noted that Columbia Slough and Smith and Bybee Lakes may provide important rearing habitat for juvenile Chinook salmon. Table B2 summarizes the CHART's PCE/watershed scores and conservation value ratings, and Figure B1 shows the overall distribution of ratings by HUC5 watershed.

Lower Columbia River Corridor

The lower Columbia River rearing and migration corridor consists of that segment of the Columbia River from the confluences of the Sandy River (Oregon) and Washougal River (Washington) to the Pacific Ocean. This corridor overlaps with the following counties: Clatsop, Columbia, and Multnomah counties in Oregon, and Clark, Cowlitz, Pacific, and Wahkiakum counties in Washington. Fish distribution and habitat use data from ODFW and WDFW identify approximately 118 miles of occupied riverine and estuarine habitat in this corridor (ODFW 2003a,b; WDFW 2003). Table B1 summarizes the total number of occupied reaches in this corridor containing rearing or migration PCEs, as well as management activities that may affect the PCEs.

After reviewing the best available scientific data for all of the areas within the freshwater and estuarine range of this ESU, the CHART concluded that the lower Columbia River corridor was of high conservation value to the ESU. Other upstream reaches of the Columbia River corridor (within the Middle Columbia/Hood and Lower Columbia/Sandy Subbasin subbasins above) are also high value for rearing/migration. The CHART noted that the lower Columbia River corridor connects every watershed and population in this ESU with the ocean and is used by rearing/migrating juveniles and migrating adults. The Columbia River estuary is a particularly important area for this ESU as both juveniles and adults make the critical physiological transition between life in freshwater and marine habitats (ISAB 200, Marriott et al. 2002).

Marine Areas

NOAA Fisheries' analysis focused on freshwater and estuarine habitats upstream of the mouth of the Columbia River. While marine areas are occupied by this ESU, within this vast area the agency has not identified "specific areas within the geographical area occupied by the species . . . on which are found those physical or biological features . . . essential to the conservation of the species."

Changes to the CHART's Initial Assessments

The CHART reviewed the public and peer reviewer comments received on the Team's initial findings for this ESU as well as new information relevant to evaluating habitat areas for this ESU. As a result, the CHART did not change conservation value ratings for any watershed within the geographical area occupied by this ESU, and there were no changes to the delineation of occupied habitat areas (although the CHART did correct a mapping error in the North Fork Toutle River that resulted in only tributaries in this HUC5 being eligible for exclusion). The proposed critical habitat designation (69 FR 74572, December 14, 2004) summarizes the comments and responses pertaining to the CHART's initial determinations for this ESU and Tables B1 and B2 reflect the final CHART assessments.

References and Sources of Information

References cited above as well as key reports and data sets reviewed by the CHART include the following:

- Bastasch, R., A. Bibao, and G. Sieglitz. 2002. Draft Willamette Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- Byrne, J. 2002a. Draft Lewis River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- Byrne, J. 2002b. Draft Washougal River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- City of Portland. 2003. Comments on NOAA Fisheries' advanced notice of proposed rulemaking. City of Portland, Oregon, Endangered Species Act Program. Letter dated November 12, 2003. (Available from NOAA Fisheries Protected Resources Division, Portland, Oregon)
- Coccoli, H. 2000. Draft Hood River Subbasin Summary (including tributaries between Bonneville Dam and the Hood River). Report Prepared for the Northwest Power

- Planning Council, dated November 15, 2000. (Available at: <http://www.cbfgwa.org/>)
- Dammers, W. 2002. Draft Cowlitz River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfgwa.org/>)
- Forest Ecosystem Management Assessment Team (FEMAT). 1993. Forest ecosystem management: an ecological, economic, and social assessment. Report of the Forest Ecosystem Management Assessment Team. U.S. Government Printing Office 1993-793-071.
- Fulton, L. 1968. Spawning areas and abundance of Chinook salmon (*Oncorhynchus tshawytscha*) in the Columbia River basin – past and present. Bureau of Commercial Fisheries Special Scientific Report – Fisheries No. 571, December 1970.
- Fulton, L. 1970. Spawning areas and abundance of steelhead trout and coho, sockeye, and chum salmon in the Columbia River basin – past and present. National Marine Fisheries Service Special Scientific Report – Fisheries No. 618, December 1970.
- Haring, D. 2003. Addendum to Wind/White Salmon Water Resource Inventory Area 29, Salmon and Steelhead Habitat Limiting Factors Analysis. Washington State Department of Ecology.
- Independent Scientific Advisory Board (ISAB). 2000. The Columbia River Estuary and the Columbia River Basin Fish and Wildlife Program. Report of the ISAB dated November 28, 2000. (Available at: <http://www.nwcouncil.org/library/isab/isab2000-5.pdf>)
- Kostow, K. (editor). 1995. Biennial Report on the Status of Wild Fish in Oregon. OR. Dep. Fish Wildl. Rep., 217 p. + app. December 1995. (Available at: <http://www.dfw.state.or.us/>)
- Lower Columbia Fish Recovery Board (LCFRB). 2003. Draft Lower Columbia Recovery Plan Technical Foundation. Working draft dated July 22, 2003. (Available from <http://www.lcfrb.gen.wa.us>)
- Lower Columbia Fish Recovery Board (LCFRB). 2004. Lower Columbia Salmon Recovery and Fish and Wildlife Subbasin Plan. December 14, 2004. (Available from <http://www.lcfrb.gen.wa.us>)

- Marriott, D., and 27 contributors. 2002. Lower Columbia River and Columbia River Estuary Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- McElhany, P., T. Backman, C. Busack, S. Heppell, S. Kolmes, A. Maule, J. Myers, D. Rawding, D. Shively, and C. Steward. 2002. Willamette/Lower Columbia Pacific salmonid viability criteria. Draft report from the Willamette/Lower Columbia Technical Recovery Team. December 2002.
- Myers, J., R. Kope, B. Bryant, D. Teel, L. Lierheimer, T. Wainwright, W. Grant, F. Waknitz, K. Neely, S. Lindley, and R. Waples. 1998. Status review of Chinook salmon from Washington, Idaho, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo NMFS-NWFSC-35, 443 p.
- Myers, J., C. Busack, D. Rawding, and A. Marshall. 2003. Historical population structure of Willamette and lower Columbia River basin Pacific salmonids. Willamette/Lower Columbia River Technical Recovery Team report. (October 2003). (Available at http://www.nwfsc.noaa.gov/trt/popid_report.htm)
- NOAA Fisheries. 2003. Preliminary conclusions regarding the updated status of listed ESUs of West Coast salmon and steelhead. Report of the West Coast Salmon Biological Review Team dated February 19, 2003.
- NMFS. 2005. Habitat Distribution for 12 Evolutionarily Significant Units of Pacific Salmon and Steelhead in Oregon, Washington, and Idaho. August 2005. GIS data available from:
<http://www.nwr.noaa.gov/1salmon/salmesa/crithab/CHsite.htm>.
- Northwest Power Planning Council. 1990. Presence/absence database from Northwest Power Planning Council's subbasin planning process. (Available at www.streamnet.org)
- Olsen, E., P. Pierce, M. McLean, and K. Hatch. 1992. Stock Summary Reports for Columbia River Anadromous Salmonids, Volume I: Oregon Subbasins Below Bonneville Dam for the Coordinated Information System. Report to Bonneville Power Administration, Contract No. 1989BP94402, Project No. 198810800, 991 electronic pages (BPA Report DOE/BP-94402-1)
- Oregon Department of Fish and Wildlife (ODFW). 1992. Clackamas River subbasin fish management plan. Portland, Oregon.

- Oregon Department of Fish and Wildlife. 2003a. Oregon Salmon and Steelhead Habitat Distribution. Natural Resources Information Management Program. (Available at: <http://rainbow.dfw.state.or.us/nrimp/>)
- Oregon Department of Fish and Wildlife. 2003b. Oregon Salmon and Steelhead Habitat Distribution at 1:24,000 Scale. Natural Resources Information Management Program. (Available at: <http://rainbow.dfw.state.or.us/nrimp/>)
- Rawding, D. 2000a. Draft Little White Salmon River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated November 15, 2000. (Available at: <http://www.cbfwa.org/>)
- Rawding, D. 2000b. Draft White Salmon River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated November 15, 2000. (Available at: <http://www.cbfwa.org/>)
- Rawding, D. 2000c. Draft Wind River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated November 15, 2000. (Available at: <http://www.cbfwa.org/>)
- Roler, R. 2000a. Draft Elochoman River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- Roler, R. 2000b. Draft Grays River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- Schroeder, R.K., K.R. Kenaston, and R.B. Lindsay. 2002. Spring Chinook salmon in the Willamette and Sandy Rivers. ODFW Annual Progress Report F-163-R-08.
- Schroeder, R.K., K.R. Kenaston, and R.B. Lindsay. 2003. Spring Chinook salmon in the Willamette and Sandy Rivers. ODFW Annual Progress Report F-163-R-08.
- Scott, W.B. and E.J. Crossman. 1973. Freshwater fishes of Canada. Fisheries Research Board of Canada, Bulletin No. 184. 966 p.
- Sieglitz, G., and 9 co-contributors. 2002. Draft Sandy Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)
- Tacoma Public Utilities. 2003. Cowlitz River Project, FERC No. 2016. Mayfield Dam Fish Guidance System Evaluations. Louver Operational Change – 2002. Report dated January 2003.

- Wade, G. 2000. Salmon and Steelhead Habitat Limiting Factors, WRIA 26 (Cowlitz). Washington State Department of Ecology.
- Wade, G. 2000. Salmon and Steelhead Habitat Limiting Factors, WRIA 27 (Lewis). Washington State Department of Ecology.
- Wade, G. 2001. Salmon and Steelhead Habitat Limiting Factors, WRIA 28 (Salmon-Washougal). Washington State Department of Ecology.
- Wade, G. 2002. Salmon and Steelhead Habitat Limiting Factors, WRIA 25 (Grays-Elochoman). Washington State Department of Ecology.
- Washington Department of Fish and Wildlife (WDFW) and Western Washington Treaty Indian Tribes (WWTIT). 1993. 1993 Washington State salmon and steelhead stock inventory (SASSI). WDFW, Olympia, WA, 212p.
- Washington Department of Fish and Wildlife (WDFW). 2003. "Fishdist: 1:24,000 (24K) and 1:100,000 (100K) Statewide Salmonid Fish Distribution". GIS data layer. (M. Hudson, data manager). Available from Washington Department of Fish and Wildlife, 600 Capitol Way N, Olympia WA 98501-1091.
- Weinheimer, J. 2002. Draft Kalama River Subbasin Summary. Report Prepared for the Northwest Power Planning Council, dated May 17, 2002. (Available at: <http://www.cbfwa.org/>)

Table B1. Summary of Occupied Areas, PCEs, and Management Activities Affecting PCEs for the Lower Columbia River Chinook Salmon ESU

Map Code	Subbasin	Watershed	Area/ Watershed (HUC5) Code	Primary Constituent Elements (PCEs)			Unoccupied but may be essential (mi)**	Management Activities***
				Spawning/ Rearing PCEs (mi)	Rearing/ Migration PCEs (mi)	Migration/ Presence PCEs (mi)*		
	Middle Columbia/ Hood	East Fork Hood River	1707010506	23.1	0	0		A, C, F, I, R
	Middle Columbia/ Hood	West Fork Hood River	1707010507	31.4	0	0		A, F, R
	Middle Columbia/ Hood	Hood River	1707010508	11.6	0	0.8		A, C, D, F, R, I, U
	Middle Columbia/ Hood	White Salmon River	1707010509	2.8	0.1	0.8	15.9 ^p	A, C, D, F, R, U
	Middle Columbia/ Hood	Little White Salmon River	1707010510	0	0	1.6		D, F, R
	Middle Columbia/ Hood	Wind River	1707010511	19	6.2	17.1		F, R, U
	Middle Columbia/ Hood	Middle Columbia/ Grays Creek	1707010512	0.6	0.1	17.1		R, U
	Middle Columbia/ Hood	Middle Columbia/ Eagle Creek	1707010513	2	0.2	10.9		D, R, U
	Lower Columbia/ Sandy	Salmon River	1708000101	15.5	0	0		F, C, R
	Lower Columbia/ Sandy	Zigzag River	1708000102	11.6	0	4		F, C, R
	Lower Columbia/ Sandy	Upper Sandy River	1708000103	12.7	0	0		F, R
	Lower Columbia/ Sandy	Middle Sandy River	1708000104	26	0.3	0		D, R, U
	Lower Columbia/ Sandy	Bull Run River	1708000105	6.5	0	0		D, F, R
	Lower Columbia/ Sandy	Washougal River	1708000106	10.9	3.6	14.3		C, F, R, S, U, W
	Lower Columbia/ Sandy	Columbia Gorge Tributaries	1708000107	6.8	10.2	27.9		C, D, F, R, U, W
	Lower Columbia/ Sandy	Lower Sandy River	1708000108	20	4.2	2.4		A, C, F, R, U
	Lower Columbia/ Sandy	Salmon Creek	1708000109	0	0	40.3		A, C, F, R, U, W
	Lewis	Upper Lewis River	1708000201	0	0	0	^q	
	Lewis	Muddy River	1708000202	0	0	0	^r	
	Lewis	Swift Reservoir	1708000203	0	0	0	^s	

^p Watershed contains unoccupied habitat above Condit Dam that may be essential for conservation.

^q The downstream dams Merwin, Yale, and Swift are barriers to fish distribution in this watershed. Unoccupied habitat areas above these dams may be essential to conservation.

^r The downstream dams Merwin, Yale, and Swift are barriers to fish distribution in this watershed. Unoccupied habitat areas above these dams may be essential to conservation.

^s Swift Dam, as well as the downstream dams Merwin and Yale, is currently a barrier to fish distribution in this watershed. Unoccupied habitat areas above these dams may be essential to conservation.

Map Code	Subbasin	Watershed	Area/ Watershed (HUC5) Code	Primary Constituent Elements (PCEs)			Unoccupied but may be essential (mi)**	Management Activities***
				Spawning/ Rearing PCEs (mi)	Rearing/ Migration PCEs (mi)	Migration/ Presence PCEs (mi)*		
	Lewis	Yale Reservoir	1708000204	0	0	0	^t	
	Lewis	East Fork Lewis River	1708000205	14.9	<0.1	7.4		A, C, F, R, S, U, W
	Lewis	Lower Lewis River	1708000206	19.2	18.3	8.1		A, C, D, F, R, U, W
	Lower Columbia/ Clatskanie	Kalama River	1708000301	40.1	0.2	13.2		C, F, R, U, W
	Lower Columbia/ Clatskanie	Beaver Creek/ Columbia River	1708000302	0	5.9	0		A, C, F, R, U, W
	Lower Columbia/ Clatskanie	Clatskanie River	1708000303	8.4	5	0		A, C, F, R, U, W
	Lower Columbia/ Clatskanie	Germany/ Abernathy	1708000304	11.5	0.1	37		A, C, F, R, U, W
	Lower Columbia/ Clatskanie	Skamokawa/ Elochoman	1708000305	11.4	0.4	26.1		A, C, F, R, W
	Lower Columbia/ Clatskanie	Plympton Creek	1708000306	1.6	7.2	0		A, C, F, R, W
	Upper Cowlitz	Headwaters Cowlitz River	1708000401	0	0	7.5		C, F, R
	Upper Cowlitz	Upper Cowlitz River	1708000402	0	0	13		C, F, R
	Upper Cowlitz	Cowlitz Valley Frontal	1708000403	0	0	34.9		A, F, R, U
	Upper Cowlitz	Upper Cispus River	1708000404	0	0	22.1		C, F, R
	Upper Cowlitz	Lower Cispus River	1708000405	0	0	26.8		C, F, R
	Lower Cowlitz	Tilton River	1708000501	0	0	24.6		C, D, F, R, U
	Lower Cowlitz	Riffe Reservoir	1708000502	0	0	30.7		A, C, D, F, R
	Lower Cowlitz	Jackson Prairie	1708000503	35.7	<0.1	21.9		A, C, D, F, R
	Lower Cowlitz	North Fork Toutle River	1708000504	0	0	0.9		F, R
	Lower Cowlitz	Green River	1708000505	26.6	0	3		F, R
	Lower Cowlitz	South Fork Toutle River	1708000506	7.6	0	17.7		F, R

^t Yale Dam, as well as downstream Merwin Dam, is currently a barrier to fish distribution in this watershed. Unoccupied habitat areas above these dams may be essential to conservation.

Map Code	Subbasin	Watershed	Area/ Watershed (HUC5) Code	Primary Constituent Elements (PCEs)			Unoccupied but may be essential (mi)**	Management Activities***
				Spawning/ Rearing PCEs (mi)	Rearing/ Migration PCEs (mi)	Migration/ Presence PCEs (mi)*		
	Lower Cowlitz	East Willapa	1708000507	8.8	0	112		A, C, F, R, U, W
	Lower Cowlitz	Coweeman	1708000508	14.3	0	46.6		A, C, F, R, U, W
	Lower Columbia	Youngs River	1708000601	15.3	28.6	0		A, C, F, I, R, U, W
	Lower Columbia	Big Creek	1708000602	9.8	6.4	0		A, C, F, I, R, W
	Lower Columbia	Grays Bay	1708000603	4.2	0.2	57.2		C, F, R, W
	Middle Willamette	Abernethy Creek	1709000704	0.6	2.6	0.1		A, C, D, R, U
	Clackamas	Eagle Creek	1709001105	13.8	3.2	0		A, F, R
	Clackamas	Lower Clackamas River	1709001106	34.8	2.7	0		A, C, D, I, R, U, W
	Lower Willamette	Johnson Creek	1709001201	1.2	8.9	0.1		A, C, I, R, U, W
	Lower Willamette	Scappoose Creek	1709001202	4.2	48.9	0		A, C, F, I, R, U, W
	Lower Willamette	Columbia Slough/ Willamette River	1709001203	0	25	0		A, C, R, U, W
	Multiple	Lower Columbia Corridor (Sandy/ Washougal to Ocean)	NA	0.1	17.4	163.9 ^u		C, D, I, R, T, U, W

* Some streams classified as “Migration/Presence PCEs” may also include rearing or spawning PCEs, but the GIS data are still undergoing review to confirm additional habitat use types.

** These watersheds contain unoccupied habitat that historically supported spawning and rearing PCEs. The CHART determined that these habitat areas/watersheds may be essential for conservation of the ESU. Since these watersheds are unoccupied, the CHART did not identify management activities.

*** This list is not exhaustive. It is intended to highlight key management activities affecting PCEs in each watershed. Activities identified are based on the general categories described by Spence et al. (1996) and summarized previously in the “Special Management Considerations or Protection” section of this report. Coding is as follows: F= forestry, G = grazing, A = agriculture, C = channel modifications/diking, R = road building/maintenance, U = urbanization, S = sand and gravel mining, M = mineral mining, D = dams, I = irrigation impoundments and withdrawals, T = river, estuary, and ocean traffic, W = wetland loss/removal, B = beaver removal, X = exotic/invasive species introductions, H = forage fish/species harvest. Primary sources for this information were the CHART and reports by LCFRB (2003), Subbasin Summary Reports of the NWPPC, and land use/land cover GIS layers from the U.S. Geological Survey.

^u The Lower Columbia River from the ocean upstream approximately 46.5 miles is considered to contain estuarine PCEs, in addition to migration and rearing (ISAB, 2000).

Table B2. Summary of Initial CHART Scores and Ratings of Conservation Value for Habitat Areas Occupied by the Lower Columbia River Chinook Salmon ESU

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Middle Columbia/Hood	East Fork Hood River	1707010506	1	2	2	1	1	3	10	Moderate HUC5 score; habitat relatively more extensive in this HUC5 than in other areas of the Gorge region; this HUC5 historically supported one of just two TRT historical spring-run populations in the Gorge region; area emphasized for supplementation efforts	High
	Middle Columbia/Hood	West Fork Hood River	1707010507	0	1	2	1	1	3	8	Moderate HUC5 score; habitat still available and this HUC5 historically supported one of just two TRT historical spring-run populations in the Gorge region; PCEs overlap with a FEMAT key watershed for at-risk anadromous salmonids	High
	Middle Columbia/Hood	Hood River	1707010508	1	1	2	1	1	3	9	Moderate HUC5 score; habitat relatively more extensive in this HUC5 than in other areas of the Gorge region; this HUC5 historically supported one of just two spring chinook populations in the Gorge region; HUC5 contains important connectivity reaches for upstream HUC5s (including one containing a FEMAT key watershed for at-risk anadromous salmonids)	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Middle Columbia/Hood	White Salmon River	1707010509	1	2	1	3	2	3	12	Moderate-high HUC5 score; PCEs limited by Condit Dam but do support a TRT fall-run core population; habitat used by nonnative spring-run fish in a watershed that historically supported one of only two spring chinook populations (including a TRT core population) in the Gorge region; Watershed contains unoccupied habitat above Condit Dam that may be essential for conservation.	High
	Middle Columbia/Hood	Little White Salmon River	1707010510	1	2	0	0	1	2	6	Low-moderate HUC5 score; limited PCEs not identified as supporting a demographically independent population, but may provide some spring-run chinook habitat that could promote conservation	Medium
	Middle Columbia/Hood	Wind River	1707010511	1	2	2	1	1	2	9	Moderate HUC5 score; habitat still available and this HUC5 supports one of four TRT historical fall-run populations (including a core population) in the Gorge region; passage over Shipherd Falls provides access to relatively extensive spring-run habitat for the Gorge region; PCEs overlap with a FEMAT key watershed for at-risk anadromous salmonids	High
	Middle Columbia/Hood	Middle Columbia/Grays Creek	1707010512	1	2	1	0	1	2	7	Moderate HUC5 score; PCEs limited in this HUC5 and likely always were due to gradient barriers and small drainage size; HUC5 supports a TRT historical core fall-run population but production likely low in this HUC5; mainstem Columbia River is high value connectivity corridor	Medium

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Middle Columbia/Hood	Middle Columbia/Eagle Creek	1707010513	1	2	1	1	1	2	8	Moderate HUC5 score; PCEs in tributary habitat in HUC5 supports two TRT historical core fall-run populations; mainstem Columbia River is high value connectivity corridor	High
	Lower Columbia/Sandy	Salmon River	1708000101	3	2	2	3	2	3	15	Highest HUC5 score for entire ESU; extensive PCEs support spring-, fall, and late fall-run populations; TRT identified spring- and late fall-runs as core and genetic legacy populations; PCEs overlap with a FEMAT key watershed for at-risk anadromous salmonids	High
	Lower Columbia/Sandy	Zigzag River	1708000102	3	2	2	3	2	3	15	Highest HUC5 score for entire ESU; extensive PCEs support spring-, fall, and late fall-run populations; TRT identified spring- and late fall-runs as core and genetic legacy populations	High
	Lower Columbia/Sandy	Upper Sandy River	1708000103	3	2	2	3	2	3	15	Highest HUC5 score for entire ESU; extensive PCEs support spring-, fall, and late fall-run populations; TRT identified spring- and late fall-runs as core and genetic legacy populations	High
	Lower Columbia/Sandy	Middle Sandy River	1708000104	3	1	2	3	2	3	14	High HUC5 score; extensive PCEs support spring-, fall, and late fall-run populations; TRT identified spring- and late fall-runs as core and genetic legacy populations; HUC5 contains important connectivity reaches for upstream HUC5s (including one containing a FEMAT key watershed for at-risk anadromous salmonids)	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Columbia/Sandy	Bull Run River	1708000105	1	1	2	3	2	3	12	Moderate-high HUC5 score; PCEs more limited due to dams in this HUC5, but still support TRT core spring- and fall-run fish; the CHART also concluded that inaccessible reaches above the Bull Run Dam complex in this HUC5 may be essential to the conservation of the ESU.	High
	Lower Columbia/Sandy	Washougal River	1708000106	1	1	2	2	1	3	10	Moderate HUC5 score; not identified as a core or genetic legacy population by TRT; other HUC5s supporting fall-run fish likely to have higher conservation value in the Cascade region	Medium
	Lower Columbia/Sandy	Columbia Gorge Tributaries	1708000107	2	2	2	2	1	2	11	Moderate-high HUC5 score; tributary habitat in HUC5 supports at least one TRT historical core fall-run population and habitat in this HUC5 likely more important for this population than the upstream HUC5; mainstem Columbia River is high value connectivity corridor supporting all upstream populations.	High
	Lower Columbia/Sandy	Lower Sandy River	1708000108	1	1	2	3	2	3	12	Moderate-high HUC5 score; PCEs support spring-, fall, and late fall-run populations; TRT identified spring- and late fall-runs as core and genetic legacy populations; important connectivity reaches for all upstream HUC5s	High
	Lower Columbia/Sandy	Salmon Creek	1708000109	1	1	1	2	1	2	8	Moderate HUC5 score; PCEs limited and degraded in this HUC5; not identified as a core population; TRT genetic legacy classification not likely attributable to fish in this HUC5; other HUC5s supporting fall-run fish likely to have higher conservation value in the Cascade region	Low

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lewis	Upper Lewis River	1708000201							*	<u>Unoccupied HUC5</u> , but population expansion into this HUC5 possibly essential for conservation; The downstream dams Merwin, Yale, and Swift are barriers to fish distribution in this watershed; Unoccupied habitat areas above these dams may be essential to conservation; nearly the entire area is a FEMAT key watershed for at-risk anadromous salmonids	Possibly High
	Lewis	Muddy River	1708000202							*	<u>Unoccupied HUC5</u> , but population expansion into this HUC5 possibly essential for conservation; The downstream dams Merwin, Yale, and Swift are barriers to fish distribution in this watershed; Unoccupied habitat areas above these dams may be essential to conservation; nearly the entire area is a FEMAT key watershed for at-risk anadromous salmonids	Possibly High
	Lewis	Swift Reservoir	1708000203							*	<u>Unoccupied HUC5</u> , but population expansion into this HUC5 possibly essential for conservation; Swift Dam, as well as downstream dams Merwin and Yale, is currently a barrier to fish distribution; Unoccupied habitat areas above these dams may be essential to conservation; HUC5 contains connectivity reaches to upstream to upstream areas that are FEMAT key watersheds for at-risk anadromous salmonids	Possibly High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lewis	Yale Reservoir	1708000204							*	Unoccupied HUC5, but population expansion into this HUC5 possibly essential for conservation; Yale Dam, as well as downstream Merwin Dam, is currently a barrier to fish distribution; Unoccupied habitat areas above these dams may be essential to conservation; HUC5 contains connectivity reaches to upstream to upstream areas that are FEMAT key watersheds for at-risk anadromous salmonids	Possibly High
	Lewis	East Fork Lewis River	1708000205	2	1	2	3	2	3	13	Moderate-high HUC5 score; PCEs support fall-run fish and TRT identified HUC5 as supporting a genetic legacy population; some of best remaining habitat of three HUC5s supporting this population; uppermost areas are a FEMAT key watershed for at-risk anadromous salmonids	High
	Lewis	Lower Lewis River	1708000206	1	1	2	3	2	3	12	Moderate-high HUC5 score; PCEs support all run types in this ESU (spring-, fall-, and late fall-run fish); TRT identified HUC5 as supporting core and genetic legacy populations; conservation of these PCEs will be especially important if historical habitats upstream are made accessible. Watershed contains unoccupied habitat areas above Merwin Dam that may be essential for conservation.	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Columbia/ Clatskanie	Kalama River	1708000301	2	1	2	1	2	3	11	Moderate-high HUC5 score; supports spring- and fall-run populations; not identified as a core or genetic legacy population by TRT; CHART uncertain of rarity/importance in this HUC5 but believed that other HUC5s may have higher conservation value in the Cascade region	Medium
	Lower Columbia/ Clatskanie	Beaver Creek/Columbia River	1708000302	1	0	1	0	1	1	4	Low-moderate HUC5 score but lowest in Coast Range region; TRT identified two historical fall-run populations in this HUC5 but present distribution limited to Clatskanie River population's historic range; PCEs are extremely limited in this HUC5 relative to others in the Coast Range region and may have very limited potential for improvement	Low
	Lower Columbia/ Clatskanie	Clatskanie River	1708000303	2	1	2	1	2	2	10	Moderate HUC5 score; PCE's support a TRT fall-run population but it is neither a core nor legacy population	Medium
	Lower Columbia/ Clatskanie	Germany/Abernathy	1708000304	2	1	2	1	1	3	10	Moderate HUC5 score; PCEs support entire range of a TRT fall-run population but it is neither a core nor legacy population; other HUC5s supporting fall-run fish likely to have higher conservation value in the Coast Range region	Medium
	Lower Columbia/ Clatskanie	Skamokawa/ Elochoman	1708000305	2	2	2	1	2	3	12	Moderate-high HUC5 score, highest in Coast Range region; PCEs support entire range of a TRT fall-run and core population	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Columbia/ Clatskanie	Plympton Creek	1708000306	1	2	2	1	2	2	10	Moderate HUC5 score; PCE's support a fall-run TRT population (but neither a core nor legacy population); may have best potential for PCE improvement of the three HUC5s supporting this population	High
	Upper Cowlitz	Headwaters Cowlitz River	1708000401	0	2	1	3	2	3	11	Moderate-high HUC5 score; PCEs support spring- and fall-run fish via trap and haul; CHART believed it was important to emphasize conservation value of upper Cowlitz/Cispus HUC5s due to their historic importance and potential to promote conservation of the ESU (i.e., Upper Cowlitz River identified by TRT as a core and genetic legacy spring-run population)	High
	Upper Cowlitz	Upper Cowlitz River	1708000402	2	1	2	3	2	3	13	Moderate-high HUC5 score; PCEs support spring- and fall-run fish via trap and haul; CHART believed it was important to emphasize conservation value of upper Cowlitz/Cispus HUC5s due to their historic importance and potential to promote conservation of the ESU (i.e., Upper Cowlitz River identified by TRT as a core and genetic legacy spring-run population); HUC5 includes a FEMAT key watershed for at-risk anadromous salmonids	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Upper Cowlitz	Cowlitz Valley Frontal	1708000403	2	1	2	3	2	3	13	Moderate-high HUC5 score; PCEs support spring- and fall-run fish via trap and haul; CHART believed it was important to emphasize conservation value of upper Cowlitz/Cispus HUC5s due to their historic importance and potential to promote conservation of the ESU (i.e., Upper Cowlitz River identified by TRT as a core and genetic legacy spring-run population)	High
	Upper Cowlitz	Upper Cispus River	1708000404	2	2	2	3	2	2	13	Moderate-high HUC5 score; PCEs support spring- and fall-run fish via trap and haul; CHART believed it was important to emphasize conservation value of upper Cowlitz/Cispus HUC5s due to their historic importance and potential to promote conservation of the ESU (i.e., Cispus River identified by TRT as a core spring-run population)	High
	Upper Cowlitz	Lower Cispus River	1708000405	2	2	2	3	2	3	14	High HUC5 score; PCEs support spring- and fall-run fish via trap and haul; CHART believed it was important to emphasize conservation value of upper Cowlitz/Cispus HUC5s due to their historic importance and potential to promote conservation of the ESU (i.e., Cispus River identified by TRT as a core spring-run population)	High
	Lower Cowlitz	Tilton River	1708000501	1	1	2	1	2	3	10	Moderate HUC5 score; PCEs support spring- and fall-run fish via trap and haul; HUC5 is only habitat for a TRT historical spring-run population	Medium

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Cowlitz	Riffe Reservoir	1708000502	1	1	1	3	2	3	11	Moderate-high HUC5 score; PCEs support spring- and fall-run fish via trap and haul; PCEs degraded due to inundation; HUC5 primarily important as rearing/migration corridor for upstream populations	High
	Lower Cowlitz	Jackson Prairie	1708000503	2	1	1	2	2	3	11	Moderate-high HUC5 score; PCEs support fall- and spring-run TRT populations (both core and legacy); some spawning PCEs in this HUC5; important as a high value rearing/migration corridor connecting upstream HUC5s/populations with the ocean	Medium
	Lower Cowlitz	North Fork Toutle River	1708000504	1	1	2	2	1	3	10	Moderate HUC5 score; PCEs support a spring- and fall-run TRT population (and the latter is a core population); PCEs in this HUC5 are very limited relative to the other three HUC5s supporting these populations; CHART noted recolonization of area despite volcano-related impacts on PCEs	Medium
	Lower Cowlitz	Green River	1708000505	2	1	2	2	2	3	12	Moderate-high HUC5 score; PCEs support a spring- and fall-run TRT population (and the latter is a core population); most of the spawning PCEs for this population may be in this HUC5; CHART noted recolonization of area despite volcano-related impacts on PCEs	High
	Lower Cowlitz	South Fork Toutle River	1708000506	2	1	2	2	2	3	12	Moderate-high HUC5 score; PCEs support a spring- and fall-run TRT population (and the latter is a core population); extensive spawning PCEs for this population in this HUC5; CHART noted recolonization of area despite volcano-related impacts on PCEs	High

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Cowlitz	East Willapa	1708000507	2	1	1	2	2	3	11	Moderate-high HUC5 score; PCEs support fall- and spring-run TRT populations (both core and legacy); some spawning PCEs and important as a high value rearing/migration corridor connecting upstream HUC5s/populations with the ocean	Medium
	Lower Cowlitz	Coweeman	1708000508	2	2	2	3	2	3	14	High HUC5 score; PCEs support a TRT genetic legacy fall-run population as well as rearing/migration for all upriver populations (fall- and spring-run) in the Cowlitz River; one of few remaining populations in ESU sustained through natural production	High
	Lower Columbia	Youngs River	1708000601	2	1	2	1	2	3	11	Moderate-high HUC5 score; PCEs support entire range of a TRT fall-run population but it is neither a core nor legacy population; CHART noted extensive releases of out-of-ESU fish in this HUC5 and believed that other HUC5s supporting fall-run fish are likely to have higher conservation value in the Coast Range region	Medium
	Lower Columbia	Big Creek	1708000602	1	2	2	1	1	3	10	Moderate HUC5 score; PCEs support entire range of a TRT fall-run and core population	High
	Lower Columbia	Grays Bay	1708000603	2	1	2	1	2	3	11	Moderate-high HUC5 score; PCEs support entire range of a TRT fall-run population but it is neither a core nor legacy population; CHART noted that relatively extensive PCEs in this HUC5 may be indicative of higher conservation value in the Coast Range region	High


Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Middle Willamette	Abernethy Creek	1709000704	1	0	1	3	1	2	8	Moderate HUC5 score; extremely limited PCEs and HUC5 not associated with a TRT population (but possibly the Clackamas River fall-run)	Low
	Clackamas	Eagle Creek	1709001105	0	2	1	1	2	2	8	Moderate HUC5 score; PCEs support a TRT fall-run and core population but are very limited in this HUC5	Low
	Clackamas	Lower Clackamas River	1709001106	3	1	2	1	2	2	11	Moderate-high HUC5 score; PCEs support a TRT fall-run and core population; this HUC5 is the primary production area for this population	High
	Lower Willamette	Johnson Creek	1709001201	1	0	2	3	2	2	10	Moderate HUC5 score; PCEs support a TRT fall-run and core population; PCE quality degraded but CHART noted that HUC5 may provide important refuge habitat for Clackamas River population and may warrant consideration for unique adaptations; Willamette River is a high value rearing/migration corridor	Medium
	Lower Willamette	Scappoose Creek	1709001202	1	1	2	1	1	3	9	Moderate HUC5 score; PCEs support at least two populations, including a TRT fall-run and core population; PCE quality degraded but CHART noted that HUC5 may provide important refuge habitat for Clackamas River population and may warrant consideration for unique adaptations; Willamette River (Multnomah Channel) is a high value rearing/migration corridor	Medium

Map Code	Subbasin	Area/ Watershed	Area/ Watershed (HUC5) Code	Scoring System (factors)						Total HUC5 Score (0-18)	Comments/ Other Considerations	CHART Rating of HUC5 Conservation Value
				1	2	3	4	5	6			
	Lower Willamette	Columbia Slough/Willamette River	1709001203	1	0	2	3	2	3	11	Moderate-high HUC5 score; PCEs support a TRT fall-run and core population and likely support rearing/migration for other Columbia River populaions in the ESU; PCE quality degraded but CHART noted that HUC5 may provide important refuge habitat for Clackamas River population and may warrant consideration for unique adaptations; Willamette River is a high value rearing/migration corridor	High
	Multiple	Lower Columbia Corridor (Sandy/ Washougal to Ocean)	NA							NS	Area not scored since many reaches are outside HUC5 boundaries. However, the CHART concluded that rearing and migration PCEs throughout this corridor are highly essential to ESU conservation	High


* Rated by CHART although HUC5 is currently blocked and unoccupied.

Figure B1. CHART Ratings of Conservation Value for Habitat Areas in HUC5 Watersheds Occupied by the Lower Columbia River Chinook Salmon ESU


Lower Columbia River Chinook
CHART Watershed Ratings



Lower Columbia Chinook Distribution
Middle Columbia/Hood Sub-basin (17070105)



Lower Columbia Chinook Distribution Lower Columbia/Sandy Sub-basin (17080001)



Map B2

Legend

▲ Cities

■ Dams

Streams (1:100,000)

Lower Columbia Chinook

Spawning/Rearing

Rearing/Migration

Migration/Presence

Watersheds

Land Ownership

Federal

Private/Other

State/Local

Tribal



Unassigned

Water

Watershed code = last two digits of sub-basin code

Note: This map is for general reference only

Lower Columbia Chinook Distribution Lewis Sub-basin (17080002)



Map B3

Legend

- Cities
- Dams
- Lakes
- Streams (1:100,000)
- Lower Columbia Chinook Spawning/Rearing
- Rearing/Migration
- Migration/Presence

Land Ownership

- Federal
- Private/Other
- State/Local
- Tribal
- Unassigned
- Water

Watersheds


0 1.5 3 6 9 12 Miles



Watershed code = last two digits of sub-basin code

Note: This map is for general reference only

Lower Columbia Chinook Distribution
Lower Columbia/Clatskanie Sub-basin (17080003)




Map B4

Legend

- Cities
- Dams
- Streams (1:100,000)
- Lower Columbia Chinook**
 - Spawning/Rearing
 - Rearing/Migration
 - Migration/Presence

Land Ownership


- Federal
- Private/Other
- State/Local
- Tribal
- Unassigned
- Water



Watershed code = last two digits of sub-basin code

Note: This map is for general reference only

Lower Columbia Chinook Distirbution Upper Cowlitz Sub-basin (17080004)




Map B5

Watershed code = last two digits of sub-basin code

Note: This map is for general reference only

Lower Columbia Chinook Distribution
Cowlitz Sub-basin (17080005)



Note: This map is for general reference only




Watershed code = last two digits of sub-basin code

Map B6

Lower Columbia Chinook Distribution
Lower Columbia Sub-basin (17080006)




Map B7




Note: This map is for general reference only

Watershed code = last two digits of sub-basin code




Lower Columbia Chinook Distribution
Middle Willamette Sub-basin (17090007)




Note: This map is for general reference only

Watershed code = last two digits of sub-basin code

Lower Columbia Chinook Distribution Clackamas Sub-basin (17090011)



Lower Columbia Chinook Distribution
Lower Willamette Sub-basin (17090012)



Map B10

Legend

- Cities
- Dams
- Lakes
- Streams (1:100,000)
- Lower Columbia Chinook Spawning/Rearing
- Lower Columbia Chinook Rearing/Migration
- Lower Columbia Chinook Migration/Presence

Land Ownership

- Federal
- Private/Other
- State/Local
- Tribal
- Unassigned
- Water

Watersheds

Watershed code = last two digits of sub-basin code

Note: This map is for general reference only